

NEWSLETTER

Belgian Pediatric COVID-19 Task Force

16/4/2020

To all Belgian Pediatricians,

*** PICU criteria and DNR codes in times of corona.**

Statement on PICU admission criteria and advanced care planning:

NL:

http://gbs-vbs.org/fileadmin/user_upload/Unions/PED/09.04.2020_Statement_Be-Pics_Covid_-_opnamecriteria.pdf

FR:

http://gbs-vbs.org/fileadmin/user_upload/Unions/PED/09.04.2020_Statement_Be-Pics_Covid_-_criteres_d_admission.pdf

*** Self care in times of corona.**

Keep on breathing, based on a statement by UZ Gent:

FR

http://gbs-vbs.org/fileadmin/user_upload/Unions/PED/AUTOSOINS_COVID19.pdf

NL

http://gbs-vbs.org/fileadmin/user_upload/Unions/PED/ZELFZORG_TIJDENS_COVID.pdf

*** FAQ update:**

FR:

https://epidmio.wiv-isp.be/ID/Documents/Covid19/COVID-19_FAQ_paeds_FR.pdf

NL:

https://epidmio.wiv-isp.be/ID/Documents/Covid19/COVID-19_FAQ_paeds_NL.pdf

New questions welcome on covidpediatrie@gmail.com



Académie belge de Pédiatrie a.d.f.

Belgische Academie voor Kindergeneeskunde f.v.

Av. de la Couronne, 20, 1050 Bruxelles – Kroonlaan, 20, 1050 Brussel – Tel: 02-649.21.47



*** What to do in times of personal protective equipment (PPE) shortage?**

FR:

http://gbs-vbs.org/fileadmin/user_upload/Unions/PED/Une_utilisation_economique_des_equipements_de_protection.pdf

NL:

http://gbs-vbs.org/fileadmin/user_upload/Unions/PED/Zuinig_omspringen_met_beschermingsmaterialen.pdf

*** Belgian Pediatric COVID Guidelines for hospitalized children (non-PICU; based on the evidence available until 31/3/2020).**

For those who didn't find them yet:

FR:

http://gbs-vbs.org/fileadmin/user_upload/Unions/PED/Belg_Recomm_COVID_ped_31mar_FR.pdf

NL:

http://gbs-vbs.org/fileadmin/user_upload/Unions/PED/Belg_Recomm_COVID_ped_31mar_NL.pdf

Tonsillar examination – infection control implications

Asymptomatically infected children

Context

Our priority is to keep ourselves and our colleagues safe whilst maintaining a pragmatic approach, and being mindful that PPE is potentially in limited supply.

Whilst the COVID-19 narrative has focused predominantly on adults, there is growing concern about the role played by asymptomatic children in the spread of infection¹. Transmission from the upper airway has been raised as a particular concern by ear, nose and throat (ENT) specialists², with viral replication shown to take place in the upper airway as well as the lower airway. This may explain why a number of paediatric and ENT healthcare professionals have developed disease in the absence of exposure to children with currently defined risk factors.

Clinical recommendations

- We recommend that the oropharynx of children should only be examined if essential.
- If the throat needs to be examined, personal protective equipment (fluid resistant surgical face mask, plastic apron and gloves) should be worn, irrespective of whether the child has symptoms consistent with COVID-19 or not.
- If a child is being tested for COVID-19, staff should routinely wear PPE (surgical face mask and apron) to collect the combined nose/throat sample.

Suspected tonsillitis in primary care or emergency departments

- During the COVID-19 pandemic, if a diagnosis of tonsillitis is suspected based on clinical history, the default becomes **not examining the throat unless absolutely necessary**.
- If using the [feverpain](#) scoring system to decide if antibiotics are indicated (validated in children 3 years and older)³, we suggest that a pragmatic approach is adopted, and automatically starting with a score of 2 in lieu of an examination seems reasonable.
- Antibiotics should be considered in children with a total feverpain score of 4 or 5 (we suggest children with a score of 3 or less receive [safety netting advice](#) alone).
- Although this is likely to result in a temporary increase in antibiotic prescribing in children, we feel that this is preferable to healthcare staff being unnecessary exposed to COVID-19. Antibiotics rarely confer a benefit in children under 3 years with tonsillitis and should only be prescribed in exceptional circumstances or if a diagnosis of scarlet fever is strongly considered.

¹ Kam KQ, Yung CF, Cui L et al. A Well Infant with Coronavirus Disease 2019 (COVID-19) with High Viral Load. *Clin Infect Dis* 2020.

² Lu D, Wang H, Yu R et al. Integrated infection control strategy to minimize nosocomial infection of corona virus disease 2019 among ENT healthcare workers. *J Hosp Infect* 2020.

³ Little P, Hobbs FDR, Moore M et al. Clinical score and rapid antigen detection test to guide antibiotic use for sore throats: randomised controlled trial of PRISM (primary care streptococcal management). 2013; **347**: f5806.



Académie belge de Pédiatrie a.d.f.
Belgische Academie voor Kindergeneeskunde f.v.
Av. de la Couronne, 20, 1050 Bruxelles – Kroonlaan, 20, 1050 Brussel – Tel: 02-649.21.47



Coordinators (for VBS/GBS and SBP/BVK):

Tyl Jonckheer & Marc Raes

Coordination scientific committee:

Dimitri Van der Linden (FR), Daan Van Brusselen (NL)

Members of the scientific committee:

Tine Boiy, Marc Hainaut, Petra Schelstraete, Julie Frère, Koen Vanden Driessche, Stefaan Van Lierde, Anne Tilmanne, Siel Daelemans, François Vermeulen, Dimitri Van der Linden, Daan Van Brusselen

Belgian Academy of Pediatrics:

Georges Casimir

VVK:

An Bael

GBPF:

Marianne Michel

European Academy of Pediatrics:

Ann De Guchtenaere

Universities:

Stephane Moniotte, Sabine Van Daele, Gunnar Buyse, Pierre Smeesters, Marie-Christine Seghaye, Yvan Vandenplas, Stijn Verhulst.

Neonatologists:

Luc Cornette

Pediatric Intensivists:

Els Duval

Observers:

Kind en Gezin: Bart Van Overmeire

ONE: Paloma Carrillo

Sciensano: Laura Cornelissen

Flemish society pediatric of nurses: Jeroen Verlinden

French speaking society pediatric nurses: Jordaan Pollet

Association Flemish Child Psychiatrists: Sofie Crommen

Belgian Pediatric Residents Association: Levi Hoste, Siel Daelemans

Spokesperson NL: Petra Schelstraete

Spokesperson FR: Dimitri Van der Linden

Secretaries (VBS/GBS): Fanny Vandamme, Raf Denayer, Aurélie François

